

TOXICOLOGY REPORT – CASE # 55269M

Peripheral blood taken from antecubital fossa (arm vein) by hospital personnel when patient was admitted at 11:45 PM on August 26, 2008.

Ethanol – 0.17G%	Opiates – Negative
Cotinine – Positive	Cannabinoids – Negative
Lidocaine – Positive	Benzodiazepines – Negative
Atropine – Positive	Metoprolol – 0.33 µg/mL
Cocaine – Trace	Caffeine – Positive
Ethylbenzoylecgonine – 0.12 µg/mL	
Benzoylecgonine – 0.36 µg/mL	
Ecgonine Methyl Ester – Positive	

Urine sample taken from patient by hospital personnel when patient was admitted at 11:45 PM on August 26, 2008.

Nicotine – Positive	Opiates – Negative
Cotinine – Positive	Cannabinoids – Negative
Lidocaine – Not Detected	Benzodiazepines – Negative
Atropine – Not Detected	Metoprolol – Positive
Cocaine – Positive	Caffeine – Positive
Ethylbenzoylecgonine – Positive	
Benzoylecgonine – Positive	
Ecgonine Methyl Ester – Positive	

TOXICOLOGY HANDBOOK

Atropine – A drug often given by emergency medical personnel to start or stimulate the heart.

Benzodiazepines – a group of prescription drugs that depress the central nervous system. They can be prescribed for insomnia, panic and anxiety, high blood pressure or to control certain types of seizures. These drugs can also be injected to control severe muscles spasms or before surgery to relax the patient. Many of these drugs can be habit forming.

Benzoylecgonine – The major metabolite of cocaine. It is biologically active but it does not contribute to the euphoria experienced by the person using cocaine. This compound does affect the coronary blood vessels and contributes to the toxicity experienced by the user.

Caffeine – A central nervous system stimulant found in coffee, tea, many soft drinks, several types of food, and many other drugs. Caffeine can increase heart rate, constrict blood vessels, and relax air passages to improve breathing or allow some voluntary muscles to contract more easily.

Cannabinoids – a group of drugs that come from the cannabis plant (marijuana). Cannabinoids increase heart rate, cause impairment of short term memory, interfere with concentration and possibly create mood changes. These compounds give the user an altered perception of time and increased feelings of hunger. Higher doses can interfere with motor skills and the ability to perform complex tasks. Individuals on cannabinoids should not operate machinery, drive or work in dangerous environments. Alcohol and cannabinoids have a synergistic effect and increase the toxicity beyond what either drug would have on its own.

Cocaine - Cocaine is a strong central nervous system stimulant that constricts blood vessels while also increasing temperature, heart rate and blood pressure. This drug may also cause feelings of restlessness, irritability and anxiety. Cocaine is powerfully addictive and a tolerance to the cocaine high may develop rapidly after first exposure. A first exposure to cocaine can cause permanent damage to the cardiac and respiratory systems. There is no "safe" dose when the drug is administered by individuals other than qualified medical personnel, as the user has no real way of knowing the concentration of the drug when it is obtained on the street. See <http://en.wikipedia.org/wiki/cocaine> and <http://www.nida.nih.gov/DrugPages/Cocaine.html>. Cocaine works by blocking the uptake of the neurotransmitters norepinephrine, serotonin and dopamine thereby producing a false sense of euphoria, or a 'high.'

Cotinine – A metabolite or byproduct that the human body creates from nicotine. It is an indicator that nicotine has been inhaled or otherwise introduced into the body.

Dopamine - Dopamine can be naturally produced by the body or added in the form of a drug. In the brain, naturally produced dopamine functions as a neurotransmitter. Dopamine can also be supplied as a drug which acts on the sympathetic central nervous system, producing effects such as increased heart rate and blood pressure. However, since dopamine administered externally cannot cross the blood-brain barrier, dopamine given as a drug does not directly affect the brain.

Ecgonine Methyl Ester – The cholinesterase enzyme converts cocaine into this substance. This metabolite of cocaine is inactive in the body.

Ethanol - A flammable, colorless chemical compound which is found in alcoholic beverages, and is also known as ethyl alcohol or grain alcohol. It is most often referred to simply as alcohol. Ethanol is a sedative hypnotic agent that can have profound impacts on judgment, fine motor function, and reaction time. If consumed excessively, ethanol can impede movement and self-protective reflexes as well as depressing respiration and heart function. For more information on how to calculate the number of drinks that typically result in a particular blood alcohol level, see <http://www.miph.org/area/ft9.html>. For further information on the effects of ethanol, see <http://www.tcada.state.tx.us/issues/danger.html>.

Ethylbenzoylecgonine – A metabolite that forms in the body when both ethanol and cocaine are present. It is also known as cocaethylene. It is biologically active, like cocaine, but takes longer to be eliminated from the body. Because of this longer elimination time, this compound is more damaging to the heart and cardiovascular system. Ethylbenzoylecgonine has similar euphoric properties to cocaine. Some studies suggest that it is more toxic than cocaine itself, particularly to the heart.

Lidocaine – a drug that can be used as a topical anesthetic or as an antiarrhythmic agent. Lidocaine is often administered in emergency situations to strengthen heart contractions.

Metoprolol – a prescription drug given for hypertension (high blood pressure). The drug works by slowing the heart rate and relaxing the blood vessels so the heart does not have to pump as hard.

Nicotine – A drug found in tobacco that typically enters the system through inhalation or absorption through lips or gums. Nicotine's effects include increases in blood pressure and heart rate, as well as increased respiration. Long term exposure to tobacco and nicotine increases the chances of cancer and heart disease and results in addiction and dependence.

Opiates – A class of drugs from the poppy plant, which includes morphine, opium and heroin. Opiates are some of the most effective pain relievers available to physicians for treating pain. However, when used improperly without medical supervision, many opiates can become addictive. Opiates affect the respiratory center in the brain and overdoses can cause a person to stop breathing.

